Mr. Rokesh Thorat<sup>1</sup> Mrs.Sudaivi Kadam<sup>2</sup>

Turkish Online Journal of Qualitative Inquiry (TOJQI) Volume 12, Issue 7, July 2021: 8002 - 8008

**Research Article** 

# Assess The Effectiveness Planned Teaching Programm On Knowledge Regarding Hypoglycemia Among The Diabetic Patient

### Mr. Rokesh Thorat<sup>1</sup> Mrs.Sudaivi Kadam<sup>2</sup>

1.Clinical Instructor "Bharti Vidyapeeth Deemed to be University, College of Nursing, Sangli,Maharashtra,India 416414.Email ID-rokesh.thorat@gmail.com

2.Clinical Instructor, Bharti Vidyapeeth Deemed to be University, College of Nursing, Sangli,Maharashtra,India 416414.Email ID-sudaivikadam@gmail.com

Corresponding Author E-mail id: rokesh.thorat@gmail.com

#### Abstract

Diabetes Mellitus (DM) is a serious global health problem. It is one of the most common metabolic disorders, with significant medical and socioeconomic implications (Wendell, et al., 1997). Diabetes mellitus is a metabolic disease with a number of causes that is characterised by chronic hypoglycemia and metabolic disturbances.

The present study was Quantitative Research approach and Pre experimental design (one group pre-test posttest) is adopted in the present study. The sample size was calculated by power analysis formula. The present study consisted of 60 patient diabetic patient. with this non probability purposive sampling technique will be used to select the sample for the present study. In the present study the samples were selected diabetic patient.

Keywords. Assess, knowledge, hypoglycemia.

#### Introduction

Diabetes mellitus is a chronic (lifelong) condition marked by high levels of glucose in the blood .According to national figures; an estimated 347 million people lived in the United States in 2008. Diabetes affected people all over the world, and the incidence was high. is rising, particularly in low- and middle-income countries. a country In India, there were 69.2 million people living with HIV/AIDS. According to 2015 results, 8.7% of people have diabetes. It is the most critical of these. remained undiagnosed for several months In 2016, India's diabetes patient population was projected to be 422 million, with the figure expected to rise to 700 million by 2030. Men have a higher prevalence of diabetes than women, but women have more diabetes than men.

#### The objectives of the study:-

- □ To assess the pre- test knowledge regarding hypoglycemia.
- $\Box$  To assess the post- test knowledge score regarding hypoglycemia.
- $\square$  To compare the pre- test and post-test knowledge score regarding hypoglycaemia.
- □ To find out the association between pre- test knowledge score and selected demographic variables

# Assess The Effectiveness Planned Teaching Programm On Knowledge Regarding Hypoglycemia Among The Diabetic Patient

### **Material and Methods**

The present study was Quantitative Research approach and Pre experimental design (one group pre-test posttest) is adopted in the present study. The sample size was calculated by power analysis formula. The present study consisted of 60 patient diabetic patient. with this non probability purposive sampling technique was used to select the sample for the present study. In the present study the samples were selected diabetic patient. The reliability was conducted at vijaynagar and wanlesswadi Dist.-Sangli of 10 samples. The reliability of the tool was done by using Karl pearson's coefficient .The 'r' value of the tool is 0.9 which found to be reliable. The pilot study was conducted in selected area Sangli, Miraj, kupwad, corporation. Dist.-Sangli at between 22-03-21 to 29-03-21.The sample size was 10. The setting and samples used for pilot study are excluded from the final study. Samples were selected by using stratified proportionate sampling technique. The purpose and significance of the study was explained to the participants . After that the data was analyzed with the help of statistician. It was revealed that the while the p value is 0.05 which is less than 0.05 and hence H1 is accepted. **Results and Discussion-**

#### Table-1

#### FREQUENCY AND PERCENTAGE DISTRIBUTION OF

#### **DEMOGRAPHIC VARIABLES**

N=60

	18	30.00
	20	33.33
	22	36.67
	30	50.00
	30	50.00
	18	30.00
	26	43.33
NDARY	6	10.00
	10	16.67
		18   26   NDARY 6   10

LINEMPL VOED	17	28.23
UNLIVIFLIOLD	17	20.33

#### TABLE NO -2

#### PRE TEST KNOWLEDGE

# FREQUENCY AND PERCENTAGE DISTRIBUTION OF SCORE OF PRE TEST KNOWLEDGE

N=60

Category	Frequency	Percentage
Poor	4	6.67%
Average	56	93.33%
Good	0	0.00

The above Table Shows That Pre Test Knowledge Score, Which Is At 6.67% Hence Diabetic Patients had Poor Knowledge Regarding the Hypoglycemia, 93.33% had Average Knowledge, and no one having Good knowledge i.e 0.00%

#### TABLE NO 3:

#### POST KNOWLEDGE

# FREQUENCY AND PERCENTAGE DISTRIBUTION OF SCORE OF POST TEST KNOWLEDGE

N=60

Category	Frequency	Percentage
Poor	0	0.00
Average	7	11.67
Good	53	88.33

Above table shows that an knowledge score, at the time of post test no one of diabetic patients had poor knowledge regarding hypoglycaemia 11.67% and 88.33% good knowledge

#### **TABLE NO: 4**

### COMPARISON OF THE MEAN AND STANDARD DEVIATION OF PRE TEST AND POST TEST KNOWLEDGE.

# Assess The Effectiveness Planned Teaching Programm On Knowledge Regarding Hypoglycemia Among The Diabetic Patient

N=60

	A. M	SD	t VALUE	p VALUE	CONCLUSION
PRE TEST	6.83	0.92	26.24	0	SIGNIFICANT
POST TEST	12.45	1.44	26.24	0	SIGNIFICANT

The Comparison of the Knowledge Score of Pre Test and Post was done By Paired t Test. The Pre Test Average Was 6.83 With Standard Deviation Of 0.92. The Post Test Average Score Was 12.45 with Standard Deviation 1.44. The Test Statistics Value Of The Paired ,,t" Test Was 26.34 With p Value 0.00. Shows That Calculated Value Is More Than Table Value There Was Significant Difference In The Average Knowledge Score , At 5% Level Of Significance.

#### TABLE: 5

### COMPARISON OF THE MEAN AND STANDARD DEVIATION OF PRE TEST AND POST TEST KNOWLEDGE.

N=60

Sr. no.	Variables	Poor	Average	Chi square	d.f.	P value	Significance	Conclusion
1	Age							
	21-30	1	17				The result	
	31-40	1	19			0.84	is not signifi	There is no association
				0.33	2	7894	cant at p <	between before planned teaching score and Age
	41 to above	2	20				.05	6
2	Gender							
							The result	There is no association

# Mr. Rokesh Thorat<sup>1</sup> Mrs.Sudaivi Kadam<sup>2</sup>

	Female	2	28				is not signifi	
				0	1	1		between before planned
							cant at p <	
	Mala	2	29				05	teaching score and Gender
	Male	2	28				.05	
3	Education							
	Primary	3	15					
							The result	There is no association
	Secondary	1	25		3			
						0.22	is not signifi	between before planned
	Higher			4.36				
						5123	cant at p <	teaching score and
	secondary		6					
							.05.	Education
	Graduate		10					
	Occupatio							
4	n							
							The result	There is no association
	Employed	2	41					
						0.31	is not signifi	between before planned
	Unemploy			0.99	1			
	e					9742	cant at p <	teaching score and
	d	2	15				.05.	Occupation

# Significant association:-

There Is No Association Between Demographical Variables.

# DISCUSSION OF THE STUDY.

The finding of the present study have been discussed as per the objectives and hypothesis of the study. The finding of the study shows that after conducting the planned teaching program there is increased in knowledge regarding hypoglycaemia among diabetic patients and statistically it was found that there is a highly significant difference in pre test and post test score.

In the present study knowledge score regarding hypoglycaemia among diabetic patients was collected by self structured questionnaire. The total score was 16and divided into 0-5 (poor), 6-10(average), 11-16 (good).

The demographic variables of diabetic patients like age, gender, education, occupation.

The mean knowledge of pre test and post test scores were evaluated and it showed significant increases in knowledge of diabetic patients who were selected for the research study and the hypothesis h1 was accepted

# **CONCLUSION:-**

The non experimental quantitative research design was used for the study, which consists of the group pre – test and post – test method . the group consists of 60 samples which were selected by simple random sampling criteria for the study, the pre –test was conducted to assess the existing knowledge about hypoglycaemia among the diabetic patients and the planned teaching programme was given on hypoglycaemia followed by post test to assess increase score in the knowledge about hypoglycaemia.

The content validity tool was done, the pilot study was conducted on 10 sample and the feasibility of the study was established from wanlesswadi sangli, conducted from 22<sup>nd</sup> march2021 to 27<sup>th</sup> march 2021.Based on the objectives and hypothesis the collected data was analysed by using descriptive and inferential statistics. Statistically mean score findings shows that planned teaching program about hypoglycaemia was effective in increasing the knowledge regarding hypoglycaemia among diabetic patients. It shows that yet diabetic patients need to gain knowledge regarding hypoglycaemia.

# **References:**

- 1. Eva T salikian, MD, (2006), Prevention of Hypoglycemia During Exercise in diabetic patients.
- 2. Implications for diabetes management and patient education (June 2007) panelDianeWild aRobynvon .
- 3. Irene S, et al. (2008) Barriers to Physical Activity among Patients with Type 1 Diabetes, knowledge and practice of nutrition related hypoglycaemia and related factors in diabetic patient. *Diabetic care* 31: 2108-2109.).
- **4.** A study conducted by Diane Wild, Robyn von .on September 2007 ,on fear of HYPOGLYCEMIA in diabetes: implications of diabetes management and patient education.
- 5. A study conducted by Timothy W Bodnar, Maria J Acevedo, Massimo Pietropaoloetal. On 2014 study published Management of Non-Islet-Cell Tumor Hypoglycemia.
- 6. A study conducted by Carrie Fidler, Torsten Elmelund Christensen, Samantha Gillard Journal of medical economics , 2011.s
- Mercede Núñez, Silvia Diaz, Tatiana Dilla ,Jesús Reviriego, Antonio Perez Diabetes Therapy 375-392, 2019...Epidemiology, quality of life, and costs associated with hypoglycemia in patients with diabetes in Spain
- 8. Patrícia BD Sanches, Isabel CF da Cruz MANAGEMENT OF HYPOGLYCEMIA
- 9. Jacqueline La Manna, Michelle L Litch man, Jane K Dickinson, Andrew Todd, Mary M
- 10. Julius, Christina R Whitehouse, Suzanne Hyer, Jan Kavookjian The diabetes educator 2019.Diabetes education impact on hypoglycemia out comes.

- 11. Erin C. Cobry and Sarah S. Jaser The Potential of Diabetes Technology to Improve Sleep in Youth With Type 1 Diabetes and Their Parents Caregivers who are responsible for the diabetes management .
- 12. Doreen Reifegerste, Sarah Hartleib Journal of clinical & translational endocrinology 2016. Hypoglycemiarelated information seeking among informal caregivers of type 2 diabetes patients
- 13. `12) Faisal K. Alanazi, BN, MN, Jazi S. Alotaibi, MN, PhD, [...], and Bander Albagawi, MN, PhD Knowledge and awareness of diabetes mellitus and its risk factors in Saudi Arabia
- 14. Faisal K. Alanazi, BN, MN, Jazi S. Alotaibi, MN, PhD, [...], and Bander Albagawi, MN, PhD Knowledge and awareness of diabetes mellitus and its risk factors in Saudi Arabia
- 15. Beth Israel Deaconess Medical Center, Boston ,Massachusetts, USA b Joslin Diabetes Center, Boston, Massachusetts, USA c Harvard Medical School, Bosto n, Massachusetts, USA Received 7 August 2015.
- 16. PE Cryer, SN Davis, <u>h Shamoon</u> Diabetes care, 2003 Am Diabetes Assoc
- 17. S Thomas, S Mathai, S Shenai... International Journal of ..., 2019 ijone.org
- 18. V Shriraam, S Mahadevan, M Anitharani... Indian journal of ..., 2015 ncbi.nlm.nih.gov