Turkish Online Journal of Qualitative Inquiry (TOJQI) Volume 12, Issue 7, July 2021: 8031 - 8036

Research Article

A study to assess the effectiveness of planned teaching program on knowledge regarding postoperative home care following coronary artery bypass grafting among patient undergoing surgery in selected hospitals.

Mr. Satish Nagkumar Salvi¹, Mrs.Nirmala Londhe²

1. Assistant Professor, M.Sc.(N) Community Health Nursing, Bharati Vidyapeeth (Deemed to be University), College of Nursing, Sangli, Maharashtra, India 416414.

Email ID: s.salavi6@gmail.com

2. Clinical Instructor, M.Sc. (N) Community Health Nursing, Bharati Vidyapeeth (Deemed to be University), College of Nursing, Sangli, Maharashtra, India 416414.

EmailID:nirmalalondhe2020@gmail.com

Corresponding Author E-mail ID: s.salavi6@gmail.com

ABSTRACT

The research design used for study was one group pre-test, post- test non experimental design to collect the sample constitute;50 samples of fourth class workers in selected private hospital. Based on the study objective pre-test and post-test was conducted and questionnaire was used for collecting the data. As the pre-test mean was 8.7 and post-test mean was 15.7, p value is 0.00001 it is highly significant as the p value is a lesser than 0.05 therefore there is improvement in knowledge score of the post-operative patients after planned teaching programme signifying that the planned teaching programme was effective. RESULT AND CONCLUSION: -There were two sections for the research study section 1 consists of demographic variable and Section 2consist of structured questionnaire. Study result shows there is significant increase in the knowledge of post of operative home care related to CABG.Researcher concluded that the planned teaching programme was effective to increase the knowledge of post of operative home care related to CABG.

Keywords: Planned teaching Programme, Coronary artery bypass grafting.

Introduction

Coronary Artery Bypass Graft (CABG) surgery, recently called aortocoronary bypass (ACB) is the surgical procedure performed to relieve angina and reduce the risk of death from coronary artery disease. Arteries or veins from elsewhere in the patient's body are grafted to the coronary arteries to bypass atherosclerotic narrowing and improve the blood supply to the coronary circulation supplying the myocardium. Commonly used conduits in CABG are the internal mammary arteries, radial arteries and the greater saphenous vein6.

During this century, the number of CABG surgeries done per day is increasing in a tremendous manner. According to American Heart Association statistics, in 2009, 5, 48,000 bypass surgeries were done in United States. Of these 3, 23, 000 were men and 2, 25,000 were women 7. Each year nearly 3, 00,000 bypass surgeries are performed in US which means 61 in every 1, 00,000 people have a

coronary bypass operation. In Britain, only 6 in every 1, 00,000 have the same operation. In Japan, 1 in 1, 00,000 patients will have a coronary bypass operation8.

A comparative study was conducted to study of the use of cardiac procedures between the United States and Ontario, Canada. 224,258 Medicare patients were compared to 9,444 patients of a similar age. In the U.S. patients 34.9% underwent coronary angiograms versus 6.7% of the Canadian patients.4For coronary artery bypass surgery, 10.6% U. S. patients underwent this treatment, versus only 1.4% of the Canadian patients. The 30 day mortality for the U.S. patients was 21.4% versus 22.3% for the Canadian patients. At one year the mortality was 34.3% in the United States and 34.4% in Canada. The study shows a strikingly higher use of cardiac procedures and the high mortality rates related to CABG.5

A retrospective survey was conducted by Escorts Heart Institute and Research Centre (EHIRC), New Delhi to explore the cardiovascular disease trends in India between 1988 and 2005. According to EHIRC data 33,686 CABG surgeries were performed in their hospitals during this period. Of these, majority (26,238) belonged to the age group 51-65 years. In 2005 alone 3500 cases were performed and the highest mortality rate (23%) was seen in patients >70 years 10.6.

Materials and Methods

Research Approach:

Research approach is the plan and procedure that consist of the steps of broad assumption to detailed method of data collection, analysis and interpretation.

In this study Quantitative research approach is used.

Research Design:

Pre-experimental one group pre-test and post-test design.

Variables under study:

Dependent variables: Knowledge regarding post-operative home care of CABG.

Independent variables: Planned teaching program

Research Setting:

The study will be conducted among selected hospitals of Sangli, Miraj and Kupwad Municipal Corporation.

Population:

Patients undergoing CABG surgery

Sample Size: 50

pic Size. 3

Sample:

Selected patients undergoing CABG surgery in selected hospitals of Sangli, Miraj Kupwad corporation area.

Sampling Criteria:

Inclusion criteria:

- **1.** Patient who are willing to participate.
- 2. Patient who can read, understand and write English and Marathi.

Exclusion criteria:

- **1.** Patient who are transferring out to another health care setting.
- 2. Patient who are planning to take DAMA.
- 3. Critically ill unresponsive clients

Sampling method:

Simple Random Sampling.

Reliability:

Reliability of the tool is done by using split half method with the help of karl Pearson's formula.

The reliability score was 0.8 which is more than 0.7 the tool was found to be reliable, feasible and acceptable. No any changes were done in tool after reliability.

Results and Discussion

SECTION I:-

FREQUENCY AND PERCENTAGE DISTRIBUTION OF DEMOGRAPHIC VARIABLES.

Table no. 1

FREQUENCY AND PERCENTAGE DISTRIBUTION OF DEMOGRAPHIC VERIABLES

N=50

Age (In years)	Frequency	Percentage
30-40	10	20.00
41-50	12	24.00
51-60	16	32.00
61-70	6	12.00
71-80	6	12.00
Gender		
Female	20	40.00
Male	30	60.00
Religion		
Hindu	20	40.00
Muslim	18	36.00
Christian	12	24.00

Education		
Primary	28	56.00
Secondary	10	20.00
Higher Secondary	12	24.00
Information Regarding		
CABG		
NO	50	100.00

TABLE NO.1 Shows that, maximum sample 32% belong to age group of 51-60 years 60% sample belong to male gender. 40% samples are Hindu. 56% sample to primary group of education

SECTION II

TABLE NO 2

COMPARISON OF PRE-TEST AND POST-TEST KNOWLEDGE SCORE ON KNOWLEDGE OF ERGONOMICS.

N=50

	A.M.	S.D.	t value	p value	Significance
Pre Test	8.62	2.36	22.2086	0	The result is
Post Test	16.38	1.23			significant at p
					< .05

Table 2 shows that the average score of post-test knowledge 16.38 was found to be significantly higher than average score of pre-test knowledge 8.62

The statistics value of the t value (table value) was 22.2086 and 'p' value 0 shows that there is significant difference in the average knowledge score, at 0.5% level of significance.

Hence, H0 is rejected as there is significant increase in the post-test knowledge score.

SECTION III -

DEALS WITH ANALYSIS OF DATA RELATED TO ASSOCIATION BETWEEN PRE- TEST KNOWLEDGE SCORE WITH SELECTED DEMOGRAPHIC VARIABLES

N = 50

Sr. no.	Variables	Chi square value	p value	Significance	Conclusion
1	Age				
	30-40	6.15	0.63	The result	No association

	41-50			is not significant	between Age and
	51-60			at $p < .05$	pre test knowledge
	61-70				score
	71-80				
2	Gender	1.88	0.39	The result	No association
	Female			is not significant	between Gender
	Male			at p $< .05$	and pre test
					knowledge score
3	Religion	0.47	0.97	The result	No association
	Hindu			is not significant	between Religion
	Muslim			at p $< .05$	and pre test
	Christian				knowledge score
4	Education	2.61	0.62	The result	No association
	Primary			is not significant	between Education
	Secondary			at p $< .05$	and pre test
	Higher				knowledge score
	secondary				

References

- 1. Kaul U, Bhatia V. Perspective on coronary interventions & cardiac surgeries in India. Indian J Med Res 132 2010 Nov; 543-48.
- 2. Medindia. Coronary Artery Bypass Grafting (CABG). [Cited 2010]. Available from www.medindia.net/Consumer Health/Health Information
- 3. Dr Alan G, Dr Marian W. Best Practice Guidelines for Cardiac Rehabilitation and Secondary Prevention [Online]. 1999 Apr 8 [cited 1999 Jun 10]; Available from: http://rubens.its.unimelb.edu.au/~heart/
- 4. 4. Baskett RJF, Buth KJ, Collicott C, Ross DB. Preoperative cardiovascular risk factor control in elective coronary artery bypass graft patients: A failure of present management. Can J Cardiol 2002;18(4): 397-402.
- 5. Ozer N. The problems and needs of caregivers after coronary artery bypass graft surgery: A literature review. 4th International Congress of Surgical and Operating Room Nursing; 2003 Oct 22-26; School of Nursing, Atatürk University, Erzurum, Turkey.
- 6. 6. Wikimedia foundation. Coronary artery bypass surgery. [cited 2010 Jun 26]. Available from http://en.wikipedia.org/wiki/Coronary_artery_bypass_surgery
- 7. American Heart Association. American heart association's heart disease and stroke statistics 2009 update [cited Nov 18 2010]. Available from www.americanheart.org/presenter.jhtml
- 8. Heart disease quotes. Heart bypass(allopathy). [cited Aug 2010]. Available from www.whale.to/m/quotes20.html
- 9. 9. 7Mortality Statistics On Bypass Surgery And Angioplasty [Cited Nov 2009]. Available from www.heartprotect.com/mortality-stats.htm
- 10. 10. Trehan N. Cardiovascular disease trends in India. Escorts Heart Institute and Research Centre; 2006 Nov 16; New Delhi, India.
- 11. 11. Suzanne C, Brenda GB, Janice LH, Kerry HC. Brunner and Suddarth's Textbook of Medical Surgical Nursing. 12th edn. Lippincott Williams & Wilkins

- 12. 12. Cesena FH, Favarato D, César DF, Oliveira SA, Luz PL. Cardiac complications during waiting for elective coronary artery bypass graft surgery: incidence, temporal distribution and predictive factors. Eur J Cardiothorac Surg 2004; 25:196-202. Available from indmed.nic.in/imcwebij.html.
- 13. 13. Lewis SM, Heitkemper MM, Dirksen SR. Medical Surgical Nursing. 6th edition. America: Mosby Publishers; 2000.
- 14. 14. Philip K, Caves. Preoperative and postoperative management of patients undergoing
- 15. 15. Coronary artery bypass graft infection. [cited Nov 2009]. Available from www.nuh.com.sg/infection/coronary-artery-bypass-graft-cabg-infections.html
- 16. 16. Bhatia JY, Pandey K, Rodrigues C, Mehta A, Joshi VR. Postoperative wound infection in patients undergoing coronary artery bypass graft surgery: a prospective study with evaluation of risk factors. Indian J Med Microbiol 2003 Oct-Dec; 21(4)246-51.
- 17. 17. Currey J, Browne J, Botti M. Haemodynamic instability after cardiac surgery: nurses' perceptions of clinical decision-making. J Clin Nurs. 2006 Sep; 15(9):1081-90.
- 18. 18. SamKG, Andrade HH, Pradhan L, PradhanA, Sones SJ, Rao PG. et al. effectiveness of an educational program to promote safety among pesticide industries to handle pesticides of south india. Int Arch occup Environ Health. (serial online). 2018 may; 81(6): 787-95. Epub 2018 oct 26.
- 19. 19. Available online URL: http://www.ncbi,nlb.nih.gov/bubmed/17962973
- 20. 20. Buranatrevedh S. Sweatsrikul P. Modle development for health promotion, occupational health hazards and accidents in pathumthani, thialanahttp;//www.ncbi.nlm.nih.gov/bubmed/? Term=a+study + on + illeffects + of + chemical + fertilizers + and ++ pesticides + on + farmer + health.
- 21. 21.Maslach C and Jackson S. (1983): The measurements of experienced burnout. Available at http://online library. Wiley. Com/doil 10,1002/ job.4030020205/pdf last accessed 11th September.