Turkish Online Journal of Qualitative Inquiry (TOJQI) Volume 12, Issue 7, July 2021: 8059 - 8066

Research Article

The aim of the study is to assess the effectiveness of planned teaching program on knowledge regarding covid-19 vaccination among staff nurses in selected hospitals.

Mrs Swati C Kurane¹ (Assistant Professor), Mr. Pratik Amrao² (Clinical Instructor)

1. Medical Surgical Nursing ,Bharati Vidyapeeth (Deemed to be University), College of Nursing, Sangli, Maharashtra, India416414

Email id: swatichandrahaskurane@gmail.com

2. Medical Surgical Nursing ,Bharati Vidyapeeth (Deemed to beUniversity) ,

College of Nursing, Sangli, Maharashtra, India416414

Email id: amraopratik55@gmail.com

ABSTRACT

The aim of the study is to assess the effectiveness of planned teaching program on knowledge regarding covid-19 vaccination among staff nurses in selected hospitals at Sangli -Miraj-Kupwad corporation area.

OBJECTIVES

1. To assess the existing knowledge regarding covid-19 vaccination.

2. To assess the effectiveness of planned teaching program on knowledge regarding covid-19 vaccination.

3. To find out the association between the pre-test knowledge score with selected demographic variable.

HYPOTHESIS

H⁰: There is no difference between the pre-test and post-test knowledge score on covid-19 vaccination.

H¹: There is difference between the pre-test and post-test knowledge score on covid-19 vaccination.

H²: There is association between pre-test knowledge score and the selected demographic variables.

METHODS : The study was conducted in Bharati hospital , Wanlesswadi ,sangli. The research design

used Pre experimental research design (One group pre-test and post-test design)

, Non-probability purposive sampling technique was used for sampling method, Sample of the study constitute of 30 Staff nurses working in Bharati Hospital , Wanlesswadi , Sangli. The tool Consists of two sections : Section-1 included 06 items seeking information on demographic profile and Section-2 Include Structured Knowledge Questionnaire on Covid-19 vaccination.

Analysis was done by using descriptive and inferential statistic. The tests used were calculation of Frequency, Percentage and paired "t" test.

RESULT: The findings of the present study showed significant difference in mean and standard deviation of pre -test and post-test. The research objective was accepted as planned teaching program is highly effective in increasing the knowledge regarding Covid-19 vaccination among staff nurses.

KEYWORDS: Effectiveness, Knowledge, Planned teaching programme, Covid-19 vaccination, Staff Nurse

INTRODUCTION:

STATEMENT OF THE PROBLEM:

"A study to assess the effectiveness of planned teaching program on knowledge regarding covid- 19 vaccination among staff nurses in selected hospitals at Sangli -Miraj-Kupwad corporation area."

OBJECTIVES OF THE STUDY:

1. To assess the existing knowledge regarding covid-19 vaccination.

2. To assess the effectiveness of planned teaching program on knowledge regarding covid-19 vaccination.

3. To find out the association between the pre-test knowledge score with selected demographic variable.

NEED FOR THE STUDY:

Coronaviruses are a big family of different viruses. Some of them cause the common cold in people which leads to severe respiratory syndrome SARS-CoV-2, the new coronavirus that causes COVID-19.Coronavirus was first detected in Wuhan, China, in late 2019 and has set off a global pandemic. COVID-19 is now having an effect across the globe. The rate of infection is growing as winter approaches.

When COVID-19 vaccines become available, the International Council of Nurses (ICN) is encouraging governments to consider the vital role of nurses and other healthcare workers by prioritizing them. The body's immune system will be stimulated after vaccination, allowing it to identify the invading pathogen and produce antibodies. This means that if the individual is exposed to the harmful pathogen again in the future, their immune system will react quickly to suppress it with antibodies, protect them from disease. The elderly and those with underlying illnesses are the most vulnerable to serious illness and death. People of all ages have become infected and have died as a result. The vaccine's efficacy in avoiding serious disease, hospitalization, and death is more significant. At this time, all three vaccines are extremely successful in avoiding COVID-19- related severe illness, hospitalization, and death.

HYPOTHESIS

H⁰: There is no difference between the pre-test and post-test knowledge score on covid-19 vaccination.

H¹: There is difference between the pre-test and post-test knowledge score on covid-19 vaccination.

H²: There is association between pre-test knowledge score and the selected demographic variables.

METHODOLOGY:

RESEARCH APPROACH :

According to polit and Hungler (1985) Research approach refers to the researchers Overall plan for obtaining answer of research questions and for testing the research hypothesis. The approach selected for the study was Quantitative research approach.

VARIABLES OF THE STUDY:

Independent variable: Planned teaching programme.

Dependent variables: Knowledge of staff nurses regarding covid-19 vaccination.

SETTING OF THE STUDY:

The place where study will be conducted. Setting for present research study is Bharati hospital & medical college Sangli.

POPULATION:

A population is an aggregate or totality of all subjects that possess a set of specification. In present study population comprises of staff nurses working in the hospitals of sangli -miraj- kupwad corporation area.

SAMPLE:

Sample is subset of population selected to participant in research study. The sample of present study consist of 30 staff nurses working in Bharati hospital and medical college sangli.

SAMPLING TECHNIQUE:

A hospital were surveyed and staff nurses were identified and listed by Non-probability purposive sampling technique.

RESEARCH DESIGN :

Research Design means how when where data is collected and analyzed. Research is important to yield maximal information with minimal information . The research design used for this study is Pre-experimental research design (one group pre- test and post- test design).

RESULTS :

Section -1 Frequency and percentage distribution of selected demographic variables.

Table 1: Frequency and percentage distribution of staff nurses according to their selected
demographic variables.n=30

		frequency	Percentage%
Age (Years)	20-29	11	36.67%
	30-39	12	40.00%
	40-49	4	13.33%
	50-59	3	10.00%
Gender	Male	13	43.33%
	Female	17	56.67%
Marital status	Married	21	70.00%
τ	Unmarried	7	23.33%
	others	2	6.67%
Education	ANM	8	26.67%
	GNM	15	50.00%
PB.B.	PB.B.sc/BSc(N)		16.67%
	M.sc(N)	2	6.67%
Work experience 1-2yrs		8	26.67%
3-4Yrs		10	33.33%
5-6Yrs		4	13.33%
7	and above	8	26.67%

Received information		
Yes	22	73.33%
No	8	26.67%
Social media	14	46.67%
Television	10	33.33%
Official Website	6	20.00%

Section-2 Frequency and Percentage distribution of knowledge score.

Table-2 Frequency and Percentage distribution of staff nurses according to their level of knowledge n=30

Sr.no						
		Pre-Test		Post-Test		
	level of knowledge	Frequency	Percentage %	Frequency	Percentage %	
1						
	Poor (0-5)	1	3.33%	0	0.00%	
2						
	Average (6-9)	26	86.67%	0	0.00%	
3						
	Good (10-13)	3	10.00%	3	10.00%	
4						
	Very Good (14-17)	0	0.00%	27	90.00%	

Above table shows that pre-test 3.33% were in poor category, 86.67% were in average category and 10.00% were in good category of knowledge. No samples were in very good. And in post-test 10.00% were in good category and only 0.00% were in average category of knowledge. No samples belongs to poor knowledge.

 Table -3 Mean and Standard deviation of pre-test and post-test knowledge of staff nurses regarding Covid-19 vaccination.

Knowledge	Mean	S.D	t value	p value
Pre-test	10.96	1.54	- 22.56	0.00001
Post-test	14.93	1.14		

The above table deals with mean and standard deviation of pre-test score are 10.96 and 1.54 respectively ,and mean and standard deviation of post-test score 14.93 and 1.14.

The 't' value of knowledge -22.56 'p'

value = 0.00001

The mean knowledge was evaluated in pre-test and post-test conducted on 30 samples and it showed significant increases in knowledge about covid-19 vaccination after conducting planned teaching programme, and the staff nurses have gained knowledge regarding covid-19 vaccination.

Section-3 Association between pre-test knowledge score and the selected demographic variables.

Table-4 Findings related to association of pre-test knowledge of staff nurses with their
demographic variables.

Demographic variables	frequency	Percentage	chi square	P value	Significance
Age (Years) 20-29	11	36.67	18	0.58	Not significant
30-39	12	40.00			
40-49	4	13.33			
50-59	3	10.00			
Gender Male	13	43.33	4.278	0.58	Not significant
Female	17	56.67			
Marital status Married	21	70.00	9.8	0.5	Not significant
Unmarried	7	23.33			
others	2	6.67			
Education ANM	8	26.67	13	0.37	Not significant
GNM	15	50.00			
PB.B.sc/BSc(N)	5	16.67			
M.sc(N)	2	6.67			
Work experience 1-2yrs	8	26.67	30.013	0.77	Not Significant
3-4Yrs	10	33.33			
5-6Yrs	4	13.33			
7 and above	8	26.67			

The above table shows that there is no significant association between demographic variables i.e Age, Gender, Marital status ,Education, Work experience with Pre-test knowledge score as the calculated 'p' value is more than 0.05

RECOMANDATIONS:

Based on the findings, the following recommendations are purposes for future research.

1) Large scale study can be conducted to generalize the findings.

2) The study related to assessment of knowledge and attitude regarding covid-19 vaccination can be done.

3) A similar study can be conducted to assess the knowledge of health team professionals about covid-19 vaccination.

CONCLUSION:

The purpose of the present study was to assess the effectiveness of planned teaching programme regarding covid-19 vaccination among staff nurses.

The pre experimental research design was used for the study, which consist of one group pretest and posttest method. The group consists of 30 sample which are selected Non-probability purposive sampling technique for the study. The pre-test was conducted to assess existing knowledge about covid-19 vaccine among staff nurses and then planned teaching programme was given on covid-19 vaccine followed by the post-test to assess increased scores in knowledge about covid-19 vaccination. The result also shows that the knowledge score of staff nurses no significant association with their demographic variables. Statistically Mean score and Standard deviation finding showed that planned teaching programme about covid-19 vaccination was effective in increasing the knowledge regarding covid-19 vaccination among staff nurses.

REFERENCES:

- Khubchandani J, Sharma S, Price JH, Wiblishauser MJ, Sharma M, Webb FJ. COVID-19 vaccination hesitancy in the United States: a rapid national assessment. Journal of Community Health. 2021 Apr;46(2):270-7.
- 2. DeRoo SS, Pudalov NJ, Fu LY. Planning for a COVID-19 vaccination program. Jama. 2020 Jun 23;323(24):2458-9.
- 3. Sherman SM, Smith LE, Sim J, Amlôt R, Cutts M, Dasch H, Rubin GJ, Sevdalis N. COVID-19 vaccination intention in the UK: results from the COVID-19 vaccination acceptability study (CoVAccS), a nationally representative cross-sectional survey. Human vaccines &immunotherapeutics. 2021 Jun 3;17(6):1612-21.
- 4. Wang J, Jing R, Lai X, Zhang H, Lyu Y, Knoll MD, Fang H. Acceptance of COVID-19 Vaccination during the COVID-19 Pandemic in China. Vaccines. 2020 Sep;8(3):482.
- Islam MS, Siddique AB, Akter R, Tasnim R, Sujan MS, Ward PR, Sikder MT. Knowledge, attitudes and perceptions towards COVID-19 vaccinations: a cross-sectional community survey in Bangladesh. medRxiv. 2021 Jan 1.
- 6. Ledda C, Costantino C, Cuccia M, Maltezou HC, Rapisarda V. Attitudes of Healthcare Personnel towards Vaccinations before and during the COVID-19 Pandemic. International Journal of Environmental Research and Public Health. 2021 Jan;18(5):2703.
- Huynh G, Van Nguyen T, Nguyen DD, Lam QM, Pham TN, Nguyen HT. Knowledge about COVID-19, beliefs and vaccination acceptance against COVID-19 among high-risk people in Ho Chi Minh City, Vietnam. Infection and Drug Resistance. 2021;14:1773.
- Kasozi KI, Laudisoit A, Osuwat LO, Batiha GE, Al Omairi NE, Aigbogun E, Ninsiima HI, Usman IM, DeTora LM, MacLeod ET, Nalugo H. A Descriptive-multivariate analysis of community knowledge, confidence, and trust in COVID-19 clinical trials among healthcare workers in Uganda. Vaccines. 2021 Mar;9(3):253.

- Li L, Wang J, Leng A, Nicholas S, Maitland E, Liu R. Will COVID-19 Vaccinations End Discrimination against COVID-19 Patients in China? New Evidence on Recovered COVID-19 Patients. Vaccines. 2021 May;9(5):490.
- 10. Motoki K, Saito T, Takano Y. Scientific literacy linked to attitudes toward COVID-19 vaccinations: A preregistered study. Frontiers in Communication. 2021;6:145.
- Badalyan AR, Hovhannisyan M, Ghavalyan G, Ter-Stepanyan MM, Cave R, Cole J, Farlow AW, Mkrtchyan H. The Knowledge and Attitude of Physicians Regarding Vaccinations in Yerevan, Armenia: Challenges for COVID-19. medRxiv. 2021 Jan 1.
- Costantino A, Topa M, Roncoroni L, Doneda L, Lombardo V, Stocco D, Gramegna A, Costantino C, Vecchi M, Elli L. COVID-19 Vaccine: A Survey of Hesitancy in Patients with Celiac Disease. Vaccines. 2021 May;9(5):511.
- Gadoth A, Halbrook M, Martin-Blais R, Gray AN, Tobin NH, Ferbas KG, Aldrovandi GM, Rimoin AW. Assessment of COVID-19 vaccine acceptance among healthcare workers in Los Angeles. Medrxiv. 2020 Jan 1.
- Gagneux-Brunon A, Detoc M, Bruel S, Tardy B, Rozaire O, Frappe P, Botelho-Nevers E. Intention to get vaccinations against COVID-19 in French healthcare workers during the first pandemic wave: a crosssectional survey. Journal of Hospital Infection. 2021 Feb 1;108:168-73.
- 15. Kwok KO, Li KK, Wei WI, Tang A, Wong SY, Lee SS. Influenza vaccine uptake, COVID-19 vaccination intention and vaccine hesitancy among nurses: A survey. International journal of nursing studies. 2021 Feb 1;114:103854.
- Riad A, Pokorná A, Attia S, Klugarová J, Koščík M, Klugar M. Prevalence of COVID-19 Vaccine Side Effects among Healthcare Workers in the Czech Republic. Journal of Clinical Medicine. 2021 Jan;10(7):1428.
- 17. Borriello A, Master D, Pellegrini A, Rose JM. Preferences for a COVID-19 vaccine in Australia. Vaccine. 2021 Jan 15;39(3):473-9.
- 18. Patrick S, Singh P, Choudhary R, Matreja PS. Acceptance and Knowledge of COVID-19 Vaccine among Health-Care Professionals in Western Uttar Pradesh, India. ActaMedica International. 2021 Jan 1;8(1):24.
- 19. Lipsitch M, Dean NE. Understanding COVID-19 vaccine efficacy. Science. 2020 Nov 13;370(6518):763-5.
- 20. Craig AM, Hughes BL, Swamy GK. COVID-19 vaccines in pregnancy. American journal of obstetrics & gynecology MFM. 2020 Dec 10:100295.