

**Ndvi Based Drought Monitoring Using Remote Sensing**

Dr.Vidhyalakshmi<sup>1</sup>, B.Sreekanth<sup>2</sup>

**ABSTRACT**

Expanding temperature and modified precipitation designs, prompts the extraordinary climate occasions like Drought which definitely influences the rural generation. Farming dry season is only the decrease in the efficiency of yields because of abnormalities in the precipitation just as abatement in the dirt dampness, which thusly influences the cost cutting of the country. As the India's agribusiness is generally subject to the Monsoon, a minor change in it influences the creation just as harvest yield radically. The rural dry spell observing, evaluation just as the board should be possible all the more precisely to the assistance of geospatial systems like Remote Sensing and GIS. The motivation behind the investigation is to dissect the vegetation worry in the locale with the computation of NDVI values. The information is utilized for the computation of NDVI. The Code of (NDVI) standardized distinction vegetation list what's more, LST, gives helpful data to agrarian dry spell checking and early admonition framework for the ranchers. By figuring the NDVI values, it tends to be plainly seen that they show a high undesirable connection. The connection among NDVI values is merely fair. The NDVI when related with the vegetation list this one tends to remain utilized in the direction distinguish the horticultural dry spell of district, as shown in this work.

**Keywords:** NDVI, Remote sensing, Vegetation index, multispectral images.

**INTRODUCTION**

Dry season may be widely portrayed as a long stretch ordinary condition of the agreement among precipitation and evapotranspiration in a particular zone, which also depends upon the advantageous start of rainstorm. In short they are rehashing atmosphere events, bringing imperative water lacks, budgetary adversities and horrible social outcomes,. According to Dracup et al. the drought is also called as "Non-attendance of precipitation as phenomenal as so since quite a while past continued impacting destructively the plant and animal existence of a spot and to deplete water supplies both for private purposes and the movement of power plants especially in those areas where precipitation is customarily sufficient for such purposes". As opposed to various kinds of genuine atmosphere or cataclysmic occasions, dry seasons regularly develop steadily and its impact varies from region to territory. Therefore the early area accept a huge activity in the alleviation system. Dry season is needed to get compound with foreseen

---

<sup>1</sup>Associate Professor, Department of Civil Engineering, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai, India

<sup>2</sup>UG Scholar, Department of Civil Engineering, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai, India.

E-mail: kvidhyakamalesh@gmail.com , sreekanthchoori@gmail.com

ecological change, additionally, the ethereal connect of drought impacted territories are foreseen to extend, which could viably influence agribusiness. Agribusiness division is by and large impacted by the start of dry season as it is significantly strong on the atmosphere, air, soil clamminess, etc. Agrarian dry season is just the decline in the productivity of harvests due to peculiarities in precipitation, increase in the temperature rate, etc., which causes a reducing in the earth clamminess. The activity of far off identifying and GIS in cultivating drought disclosure, evaluation and the board is ending up being fundamental these days as they give ground breaking information in different extent of 3-D and common measures which is wild and time eating up when done by standard methodologies, for instance, Field Survey, and analyzing surveys. The vegetation record has been considered by different analysts as one of the huge boundaries for the planning of cultivating fields, estimating atmosphere impacts, processing biomass, crop yield, dry season conditions and choosing the intensity of the vegetation. NDVI was first suggested by Tucker in 1979 as a record of vegetation prosperity and thickness. Using the NDVI data of the area, the changes in plant life spread

Present-day in the domain and moreover the example in occasion of plant dry season can be considered. This record isn't freed from disfigurements, for instance, data botch during stormy season, drenching sway on thick vegetation, etc. So it is for each situation better to mix it with various boundaries to ensure more precision. It is perceived that close by exists a strong association among exterior high temperature and NDVI. It a tolerable marker of the essentialness stability at the Earth's external which can give huge information almost the apparent physical properties and air. GNDVI saw a large a result because of changes in vegetation spread and soil sogginess, and arraigned than that the exterior temperature can increase rapidly with liquid pressure. Along it might be seen that the extent of NDVI increases during many seasons of dry season. This assessment base on the plant dry season evaluation of the Raichur district through the examining of vegetation stress achieved by the minor precipitation, advanced temperature, etc., using the MODIS induced multi short lived data of two biophysical boundaries specifically, NDVI and LST.

### NDVI

Normalised Difference Vegetation Index (NDVI) measures the amount of greenness of a particular area in a region. NDVI depends on the rule that effectively developing green plants firmly retain radiation in the noticeable area of the range "photosynthetically dynamic radiation"), while unequivocally reflecting radiation in the close infrared locale. The Green-Seeker Handheld Optical Sensor Unit is an instrument for crop investigate and counselling and gives exact estimation and information logging of the NDVI. A particular region retention and impression of photograph artificially dynamic radiation over a given timeframe can be utilized to describe the wellbeing of the vegetation there, comparative with the standard with the count of NDVI formula

$$NDVI = (\lambda_{NIR} - \lambda_{RED}) / (\lambda_{NIR} + \lambda_{RED})$$

### METHODOLOGY

The NDVI procedure is mainly used for separating the various highlights introduced among the 3-band Satellite. Vegetation Cover is said to be the one of most significant biophysical pointer to topsoil disintegration, which can be there to assess and utilize the vegetation records got from the

## NDVI BASED DROUGHT MONITORING USING REMOTE SENSING

Satellite images. Vegetation parameters which enable us to depict the circulation of plant life and soil dependent scheduled the trademark reflectance examples of green vegetation. The NDVI is a basic arithmetic pointer that can be utilized to break down the satellite remote detecting observations, since a remote stage and evaluate whether the objective or article being observed which contains live green vegetation or not, NDVI is determined a


$$RNDVI = \frac{NIR - RED}{NIR + RED} \quad \text{where } (0 < NDVI < 1)$$

$$GNDVI = \frac{NIR - GREEN}{NIR + GREEN} \quad \text{where } (0 < NDVI < 1)$$

RED Wave Length is said to be visible red reflectance, and on other Wave length, NIR is close to infrared reflectance. The Band width of the NIR band is said to be between 750-1300 nm, and Red wave length has bandwidth of 600-700 nm, and where as Green band has 550 nm. The NDVI is propelled by green vegetation, which has the distinction between the two bands which are NIR and red band; it said to be higher for mansignificant chlorophyll thickness. It needs the (NIR - red) contrast and standardizes it to change the effects of lopsided enlightenment, for example, shades of hazes or slopes

### FLOW CHART OF METHODOLOGY

The level of greenness has equivalent to that of the chlorophyll focus. NDVI standards change through the ingestion of red light which has fallen on plant chlorophyll then the impression of infrared radiation by water-filled greenery cells. Entirely obvious reaches are caught by the Satellite images in type of groups concluded which highlights can be removed later applying the NDVI strategy for various qualities. The groups are communicated as far as wavelengths, in the request for 1  $\mu$ m. In spite of the fact that, the other highlights can be removed with the assistance of these four unmistakable extents (obvious blue, centre infrared, warm infrared also, centre infrared), just three noticeable groups are utilized (close to infrared, unmistakable red and obvious green) in this effort for the element extraction.



**Fig. Flowchart of Methodology**

**TO FIND VEGETATION THROUGH THE MULTI SPECTRAL IMAGES:-**

NDVI procedure requires isolating every single band, for the identification of the vegetative record through a Multi Spectral Remote Sensing image which is available in the Satellite image. When the diverse band are isolated, NDVI strategy is signed by its trademark like vegetative greenness at various NDVI limit esteems, for example, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4 and 0.5. Different NDVI limit esteems are being utilized to separate best outcome from Satellite imagery.

Band	Name of the band	Wavelength (µm)	Characteristics and usage
1.	Visible blue	0.45 - 0.52	Maximum Water Penetration
2.	Visible Green	0.52 - 0.60	Good for measuring plant vigor
3.	Visible Red	0.63 - 0.69	Vegetation Discrimination
4.	Near InfraRed	0.76 - 0.90	Biomass and shoreline mapping
5.	Middle InfraRed	1.55 - 1.756	Moisture Content of Soil
6.	Thermal Infrared	10.4 – 12.5	Soil moisture and Thermal Mapping
7.	Middle InfraRed	2.08-2.35	Mineral mapping

**RESULTS AND DISCUSSION**

The NDVI have been utilized broadly to analyze the connection between Spectral inconstancy and the adjustments in vegetation development rate. It is additionally helpful to decide the creation of green vegetation just as recognize vegetation changes. The outcomes, speak to the

## NDVI BASED DROUGHT MONITORING USING REMOTE SENSING

different highlights, which have been extricated from the satellite picture . The outcome is contrasted and the Google earth picture, which shows the improvement essentially.

Satellite remote detecting innovation is generally utilized for checking crops and agrarian dry season assessment. Different vegetation records are accessible today, however none of the significant files is viewed as characteristically better than the rest in all conditions, some files are more qualified than others for specific employments. NDVI because of its basic count is to a great extent utilized for the vegetation considers in a territorial just as worldwide level. It is constantly prudent to combine the NDVI alongside different parameters to show signs of improvement results. The LST when associated with the vegetation index it very well may be utilized to identify the horticultural dry spell of an area, as exhibited in this work

### REFERENCES

1. Arshad S, Morid S, Reza Mobasheri.M., and Agha Alikhani.M. (2008). Development of Agricultural Drought Risk Assessment Model for Kermanshah Province (Iran), using satellite data and intelligence methods. Option Meditterrianeennes, Series A, No:80.
2. Brian D.Wardlow, Martha C. Anderson, James P. Verdin (2012). Remote Sensing of Drought.Taylor& Francis Group.
3. Chakraborty, A., Sehgal, V.K., (2010), Assessment of Agricultural Drought Using MODIS Derived Normalized Difference Water Index, Journal of Agricultural Physics, Vol. 10, pp. 28-36.
4. Dabrowska-Zielinska K., Kogan F., Ciolkosz A., Gruszczynska M. &Kowalik W. (2002): Modelling of crop growth conditions and crop yield in Poland using AVHRRbased indices, International Journal of Remote Sensing, 23:6, 1109-1123.
5. Dracup, J.A., Lee, K.S. and Paulson, E.G. Jr., (1980).On the Definition of Droughts. Water Resources Research, 16 (2): 297–302. Goetz, S. J., 1997.Multi-sensor analysis of NDVI, surface temperature and biophysical variables at a mixed grassland site. International Journal of Remote Sensing, 18, 71–94. Hasan Murad and A. K. M. Saiful Islam, (2011), Drought Assessment Using Remote Sensing And Gis In North-West Region Of Bangladesh, 3rd International Conference on Water & Flood Management (ICWFM-2011).
6. Hielkema, J. U., Prince, S. D., &Aste, W. L. (1986). Rainfall and vegetation monitoring in the Savanna zone of Democratic Republic Sudan using NOAA Advanced Very High Resolution Radiometer. International Journal of Remote Sensing, 7, 1499–1514. IPCC. (2007).
7. Climate Change 2007: Synthesis Report.Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. (Core Writing Team, Pachauri R K, Reisinger A, eds.). IPCC, Geneva, Switzerland.p. 104. Jiang Zhu, Amy E. Miller, Chuck Lindsay, DayneBroderson,
8. Tom Heinrichs, Parker Martyn, (2013). ModisNdvI Products and Metrics User Manual, Version 1.0.. Justice, C.O.& Townshend, J.R.G.&Holben, B.N.& Tucker, C.J., (1985), Analysis of the phenology of global vegetation using meteorological satellite data.
9. International Journal of Remote Sensing, 6, 1271–1318. Kogan, F. N., (1987a), Vegetation index for areal analysis of crop conditions. Proceedingsof 18th Conference on Agricultural and Forest Meteorology, AMS, W. Lafayette, Indiana, on 15–18 September 1987 (Indiana, USA), pp. 103-106.
10. Thenkabail, P. S., Gamage, M. S. D. N. and Smakhtin, V. U., (2004). The Use of Remote-Sensing Data for Drought Assessment and Monitoring in Southwest Asia

11. Ahmadi H, Nusrath A, "Vegetation change Detection of Neka river in Iran by using remotesensing and GIS", *Journal of geography and Geology*, 2 (1)., pp. 58-67., 2012
12. Karaburun A. A. K. Bhandari, "Estimation of C factor for soil erosion modelling using NDVI in Buyukcekmece watershed", *Ozean Journal of applied sciences* 3, 77-85., 2010.
13. Chouhan R, Rao N," Vegetation detection in Multi spectral remote sensing images: protective Role-analysis of coastal vegetation in 2004 Indian Ocean Tsunami. *Geo-Information for Disaster Management* ", *Procedia Technology* 6, pp., 612 –621., 2012
14. Xie Y, Zhao X, Li L, Wang H," Calculating NDVI for landsat7-etm data after atmospheric Correction using 6s model: a case study in Zhangye City, China, In proceeding *IEEE Geo Informatics 18th international conference*, pp. 1-4., 2010.
15. Hu Y, Ban Y, Zhang X, Liu J, Zhuang. D," Spatial- temporal pattern of GIMMS NDVI and its dynamics in Mongolian Plateau", In *IEEE Proceeding on earth observation and remotesensing applications*, pp.1-6., 2008.
16. Merwan Achibeta, Stefan Balevb, Antoine Dutotb, Damien Olivierb," A Model of Road Network and Buildings Extension Co-Evolution *Agent-based Modeling and Simulation of Cities Procedia Computer Science*", 32, pp. 828 –833. 2014.
17. Shikhar Deep , Akansha Saklani, " Urban Sprawl modeling using cellular automata" , *The Egyptian Journal of Remote Sensing and Space Sciences*, 17, pp. 179–18., 2014.
18. A.K. Bhandari, A. Kumar, "Feature Extraction using Normalized Difference Vegetation Index (NDVI): A Case Study of Jabalpur City", *Proceedings of Communication, Computing & Security. Procedia Technology Volume 6*, pp. 612–621, 2012
19. Nageswara PPR, Shobha SV, Ramesh, KS, Somashekhar RK," Satellite -based assessment of Agricultural drought in Karnataka State, *Journal of the Indian society of remote sensing* ", 33 (3), pp. 429-434., 2005
20. Yang Y, Zhu J, Zhao C, Liu S, Tong, The spatial continuity study of NDVI based on Kriging and BPNN algorithm" , *Journal of Mathematical and computer modelling.*, pp. 77 - 85, 2010.