

Association Of Age And Gender Prevalence In Patients Undergoing Aesthetic Corrections Using Veneers

Sahil Choudhari,

Saveetha Dental College And Hospitals,
Saveetha Institute Of Medical And Technical Sciences (Simats),
Saveetha University,
Chennai, India.
Email Id: 151501027.Sdc@Saveetha.Com

Subash Sharma,

Reader,
Department Of Conservative Dentistry And Endodontics,
Saveetha Dental College And Hospitals,
Saveetha Institute Of Medical And Technical Sciences (Simats),
Saveetha University,
Chennai, India
Email Id: Subash@Saveetha.Com

Jaiganesh Ramamurthy,

Professor And Head,
Department Of Periodontics,
Saveetha Dental College And Hospitals,
Saveetha Institute Of Medical And Technical Sciences (Simats),
Saveetha University,
Chennai, India
Email Id: Jaiganeshr@Saveetha.Com

Abstract:

Aim: The Aim Of The Study Was To Find Out The Association Between The Age And Gender Distribution In Patients Undergoing Veneering Treatment.

Materials And Methods: Patients Who Reported For Esthetic Management Of Teeth And For Whom Veneering Treatment Had Been Done In Saveetha Dental College And Hospitals, Chennai From June 2019-March 2020 Were Recruited. A Total Of 156 Direct And Indirect Veneers Were Done Among The Reported Patients. Data Regarding Age, Gender, Tooth Number And Reason (Spacing, Discolouration Or Both) For Need Of Esthetic Treatment Were Collected And Analyzed.

Results: Out Of 156 Patients, Most (71.2%) Of The Patients Belonged To The Age Group Between 20 To 40 Years. Gender Distribution Of Patients Showed 37.8% Of The Patients Were Females And 62.2% Of The Patients Were Males. 48.7% Of The Patients Reported With The Complaint Of Spacing Also, Association Between Gender And Reason For Veneering Treatment And Age And Reason For Veneering Treatment Were Found To Be Statistically Significant.

Conclusion: Within The Limitations Of The Study, The Age Group Between 20-40 Years Reported The Maximum For Veneering Treatment (71.2%). Among The Gender, Males (62.2%) Reported With A Higher Incidence Than Females (37.8). Spacing (48.7%) Was The Most Common Reason For Patients Undergoing Veneering Treatment.

Keywords: Discolouration; Spacing; Veneering.

Introduction:

While The Invention Of Veneering Anterior Teeth By Dr Pincus [1] Was Presented In 1937, It Became More Popular In The Mid-1970s, Using Three Different Approaches: Direct Bonding Using Resin Composites, Prefabricated Composite Veneers, And Indirect, Custom-Made Porcelain Veneers. [2-4] In The Past Few Years, Patient Requests For Esthetic Treatment Of Anterior Teeth Have Increased. [5]

Dental Caries Is The Most Common Cause For The Loss Of Enamel In A Clinical Situation. Dental Caries Are Easily Detectable And Reversible At An Early Stage. [6] Many Clinical Situations Such As Tooth Discoloration, Extensive Fractures, Malaligned Teeth Or Dental Caries Lesions May Cause An Important Impairment In Esthetic Appearance And Smile Harmony, Causing Impact In The Quality Of Life. [7,8] Dental Trauma Is One Of The Most Commonly Seen Injuries In The General Population Involving Teeth And Other Surrounding Structures. [9] If A Patient Only Reports With Chipped Teeth Or Localised Defects, Veneers Are Usually The Material Of Choice Due To A Conservative, Esthetic Approach As They Are Tooth Coloured Restorations In Order To Give The Patient A Perfect Smile. [10] The Use Of Veneers May Be An Interesting Option To Restore The Esthetic Appearance Of Damaged Teeth, [11] Especially Because Indirect Techniques Require More Removal Of Sound Tooth Structure And Have A Higher Cost, Due To The Laboratory Procedures Involved. [12]

Universally, The Following Clinical Situations Are Considered Indications For Anterior Restorations: 1) Enamel Hypoplasia, Waves, Stains, Grooves, Etc, 2) Enamel Abrasions, 3) Congenital Amelogenesis Imperfecta, 4) Caused By Hormones Or Tetracyclines, 5) Chromatic Or Dystrophic Alterations Caused By Fluorosis, 6) Numerous Esthetically Unsatisfactory Superficial Restorations, 7) Coronal Fractures Located Primarily On The Palatal Side, 8) Retention Of Deciduous Teeth With A Resorbed Root, 8) Lateral Incisor Agensis, When The Canine Has Transposed To That Position, 9) Volume Anomalies (Microdens), 10) Diastemas, And 11) Alignment Defects [13]

Diagnosing The Exact Pulpal Status By Direct Examination Is Uncertain Due To The Fact That The Pulp Is Enclosed Within A Hard Tissue. In Order To Identify The Actual Pulp Status Surrogate Test Must Be Performed. [14] These Tests Must Be Performed Before Planning For A Veneer. Restorative Aesthetic Dentistry Should Be Practiced As Conservatively As Possible. Currently, The Use Of Adhesive Techniques Makes It Possible To Conserve As Much Tooth Structure As Is Possible While Satisfying The Patient's Restorative Demands And Aesthetic Needs. With Indirect Restorations, Clinicians Should Choose A Material And Technique That Allows The Most Conservative Treatment; Satisfies The Patient's Aesthetic, Structural, And Biologic Requirements, And Has The Mechanical Requirements To Provide Clinical Durability. [15]

Today, This Evolution Is Clinically Represented By The Choice Of Direct And Indirect Techniques. Direct Techniques Are One Session Procedures Performed Chairside By Directly Applying Resin Composite To The Dental Surface. They Are Used For Simple Restorations Using An Anatomic Layering Procedure, Which Aids The Clinician In Correctly Defining The Color And Shape Of The Tooth, Using The Residual Dental Structure As Reference. [16]

Indirect Techniques Demand At Least Two Sessions And The Collaboration Of A Dental Technician, Who Will Manufacture A Veneer To Be Luted To The Prepared Dental Surface. Indirect Techniques Are Preferred In More Complex Cases, In Which Restoring Harmonious Tooth Shape And Color Is Highly Dependent On Variables Such As The Clinician's Skill And The Technique And Material Used. Further, It Should Be Considered Whether The Patient Will More Or Less Be Compliant About A Prolonged Session. [17,18] The Advantages Offered By Indirect Techniques Compared To Direct Techniques Are As Follows: [19] Superior Esthetic Result, Adequate Abrasion Resistance, Biocompatibility With Soft Tissues, Dimensional And Chromatic Stability Over Time And Strong Bond Between The Two Adhesive Interfaces (Luting Agent/Etched Enamel And Luting Agent/Etched Porcelain Or Post-Polymerized Resin Composite). Previously Our Team Has A Rich Experience In Working On Various Research Projects Across Multiple Disciplines The [20-22][23-34].

The Aim Of This Study Was To Find Out The Association Between The Age And Gender Distribution In Patients Undergoing Veneering Treatment. It Also Highlighted The Reason Why The Patients Had Come.

Materials And Methods:

The Study Setting For This Study Was A University Setting. Patients Who Reported For Esthetic Management Of Teeth And For Whom Veneering Treatment Had Been Done In Saveetha Dental College And

Hospitals, Chennai From June 2019 To March 2020 Were Recruited In Our Study. A Total Of 156 Direct And Indirect Veneers Were Done Among The Patients Reported. Cross Verification Of Data Was Done Using Photographs. Data Was Reviewed By An External Reviewer.

Data Collected Included Name, Age, Gender, Tooth Number And Reason (Spacing, Discolouration Or Both) For Need Of Esthetic Treatment. The Collected Data Was Tabulated Using Microsoft Excel And Analysed Using Spss. Differential (Frequency Distribution And Percentage) And Inferential (Chi-Square Test) Statistics Were Done.

Results:

Our Study Included 156 Veneering Treatment Procedures Done Between June 2019 To March 2020. The Ages Were Grouped Into 4 Categories: Below 20 Years, Between 20 To 40 Years, Between 40 To 60 Years And Above 60 Years. 11.5% Of The Patients Belonged To The Age Group Of Below 20 Years, 71.2% Of The Patients Belonged To The Age Group Between 20 To 40 Years, 5.8% Belonged To The Age Group Between 40 To 60 Years And 11.5% Of The Patients Belonged To The Age Group Of Above 60 Years. (Table 1)

Gender Distribution Of Patients Showed 37.8% Of The Patients Were Females And 62.2% Of The Patients Were Males. (Table 2)

Also, The Reason For Undergoing Veneering Treatment Was Recorded. 41% Of The Patients Reported With The Complaint Of Discolouration, 48.7% Of The Patients Reported With The Complaint Of Spacing And 10.3% Of The Patients Reported With The Complaint Of Both Spacing And Discolouration. (Table 3)

Association Between Age Group And Reason For Veneering Treatment Showed That In The Age Group Of Below 20 Years, 5.13% Reported With A Complaint Of Discolouration And 6.41% Of The Patients Reported With A Complaint Of Spacing. In The Age Group Between 20 To 40 Years, 24.36% Reported With A Complaint Of Discolouration, 36.54% Reported With A Complaint Of Spacing And 10.26% Of The Patients Reported With Complaints Of Both Spacing And Discolouration. In The Age Group Between 40-60 Years, Spacing Was The Only Complaint (5.77%) And In The Age Group Of Above 60 Years, Discolouration Was The Only Complaint. (11.54%) Chi-Square Test Was Done And The Association Was Found To Be Statistically Significant. ($P = 0.00$) (Figure 1)

Association Between Gender And Reason For Veneering Treatment Showed That Among Females, 10.26% Of The Patients Reported With A Complaint Of Discolouration And 27.56% Of The Patients Reported With A Complaint Of Spacing. Among Males, 30.77% Reported With A Complaint Of Discoloration, 21.15% Reported With A Complaint Of Spacing And 10.26% Reported With Complaints Of Both Spacing And Discoloration. Chi-Square Test Was Done And The Association Was Found To Be Statistically Significant. ($P=0.00$) (Figure 2)

Discussion:

This Study Assessed The Association Of Age And Gender With Veneering Treatment. The Reason For Veneering Treatment Also Was Taken Into Consideration. Age Group Between 20 To 40 Years Reported The Maximum For Veneering Treatment (71.2%) And The Age Group Between 40 To 60 Years Reported The Least For Veneering Treatment. (5.8%) Males (62.2%) Reported With A Higher Incidence For Veneering Treatment Than Females. (37.8%)

Taking The Reason Into Consideration, Spacing (48.7%) Was The Most Common Reason For Patients Choosing Veneering Treatment Followed By Discolouration. (41%) Only 10.3% Of Veneers Fabricated Were For Both Discolouration And Spacing. In The Age Group Between 40-60 Years, All The Patients Reported With The Complaint Of Spacing And All Patients Above 60 Years Reported With The Complaint Of Discoloration.

In A Retrospective Study Done By Coelho Et Al On Direct Anterior Composite Veneers In Vital And Non-Vital Teeth, [35] 196 Restorations Were Evaluated In 86 Adult Patients (Mean Age 44 Years Old), With An Overall Success Rate Of 80.1%. The Majority Of Patients Were Females (69.8%). 83.2% Of Veneers Were Placed In Upper Central Incisors. Almost 80% Of The Evaluated Veneers Were Performed Using Universal Composites And 73% Of The Restorations Were Placed In Vital Teeth.

Study Done By Meijering Et Al On Patient's Satisfaction With Different Types Of Veneer Restorations, [36] Highlighted The Reasons For Veneer Treatment. The Reason 'Shape' And 'Position' Were More Important For The Dentist Than For The Patient, While The Reason 'Colour' Was More Important For The Patient.

Study Done By Burke Et Al On Ten-Year Outcome Of Porcelain Laminate Veneers Placed Within The General Dental Services In England And Wales, [37] 1177 Different Adult Patients Were Identified Who Had Received Porcelain Veneers, Of Whom 410 (35%) Were Males And 767 (65%) Were Females. A Number Of Patient Factors Were Found To Influence Veneer Survival. Veneers Perform Least Well In Patients Over The Age Of 60, With Reasons For This Potentially Being Deteriorating Periodontal Condition, Increased Loading Because Of

Reduced Posterior Support, Or Reduced Salivary Flow As A Result Of The Prescription Of A Variety Of Medications. Similar Findings Are Apparent For Directly Placed Restorations. [38]

Study By Dunne Et Al On Clinical Performance Of Porcelain Veneers, Age, Gender, Fabrication Technique (Platinum Foil Or Refractory Die), Use Of Rubber Dam And Year Of Bonding Were Not Significant Factors. [39]

The Limitations Of Our Study Were That It Was An Institutional Based Study, The Duration Of Cases Taken Into Account Was Only 1 Year And Limited Sample Size. Our Institution Is Passionate About High Quality Evidence Based Research And Has Excelled In Various Fields ([40–50]. Future Scope Includes Taking Larger Populations Into Account, Populations From Different Geographical Variations, Checking For Success Versus Failure Rates And Patient Satisfaction After Veneering Procedures.

Conclusion:

Within The Limitations Of The Study, Patients Between The Age Group Of 20-40 Years (71.2%) Reported To The Maximum For Veneering Treatment. Among The Gender, Males (62.2%) Reported With A Higher Incidence Than Females (37.8%). Also, Spacing (48.7%) Was The Most Common Reason For Patients Undergoing Veneering Treatment.

Acknowledgment:

The Authors Of This Study Would Like To Express Their Gratitude Towards Everyone Who Facilitated And Enabled Us To Carry Out This Study Successfully. We Would Also Like To Thank The Institute For Helping Us To Have Access To All The Case Records For Collecting The Required Cases For Conducting This Study.

Author Contribution:

Sahil Choudhari Carried Out The Retrospective Study By Collecting Data And Drafted The Manuscript After Performing The Statistical Analysis. Subash Sharma Aided In The Conception Of The Topic, Participated In The Study Design, Statistical Analysis And Supervised In Preparation Of The Manuscript. Jaiganesh Ramamurthy Participated In The Study Design And Coordinated In Developing The Manuscript. All The Authors Had Equally Contributed In Developing The Manuscript.

Conflict Of Interest:

There Was No Conflict Of Interest As Declared By The Authors.

References:

1. Pincus, R C. Building Mouth Personality. J South Calif Dent Assoc. 1938;14:125–9.
2. Faunce Fr, Myers Dr. Laminate Veneer Restoration Of Permanent Incisors. J Am Dent Assoc. 1976;93(4):790–2.
3. Helpin Ml, Fleming Je. Laboratory Technique For The Laminate Veneer Restoration. Pediatr Dent. 1982;4(1):48–50.
4. Haas Br. Mastique Veneers: A Cosmetic And Financial Alternative In Post-Periodontal Care. J N J Dent Assoc. 1982;53(4):25.
5. Pippin Dj, Mixson Jm, Soldan-Els Ap. Clinical Evaluation Of Restored Maxillary Incisors: Veneers Vs. Pfm Crowns. The Journal Of The American Dental Association. 1995 Nov 1;126(11):1523–9.
6. Rajendran R, Kunjusankaran Rn, Sandhya R, Anilkumar A, Santhosh R, Patil Sr. Comparative Evaluation Of Remineralizing Potential Of A Paste Containing Bioactive Glass And A Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An In Vitro Study [Internet]. Vol. 19, Pesquisa Brasileira Em Odontopediatria E Clínica Integrada. 2019. P. 1–10. Available From: [Http://Dx.Doi.Org/10.4034/Pboci.2019.191.61](http://Dx.Doi.Org/10.4034/Pboci.2019.191.61)
7. Sadowsky Sj. An Overview Of Treatment Considerations For Esthetic Restorations: A Review Of The Literature. J Prosthet Dent. 2006 Dec 1;96(6):433–42.
8. Meyenberg K. The Ideal Restoration Of Endodontically Treated Teeth--Structural And Esthetic Considerations: A Review Of The Literature And Clinical Guidelines For The Restorative Clinician. Eur J Esthet Dent. 2013;8(2):238–68.

9. Jose J, Subbaiyan H. Different Treatment Modalities Followed By Dental Practitioners For Ellis Class 2 Fracture–A Questionnaire-Based Survey. *Open Dent J* [Internet]. 2020; Available From: <https://Opdentistryjournal.Com/Volume/14/Page/59/Fulltext/>
10. Ravinthar K, Others. Recent Advancements In Laminates And Veneers In Dentistry. *Research Journal Of Pharmacy And Technology*. 2018;11(2):785–7.
11. Prieto Lt, Araujo Ct, De Oliveira Dc, De Azevedo Vaz Sl, D'arce Mb, Paulillo La. Minimally Invasive Cosmetic Dentistry: Smile Reconstruction Using Direct Resin Bonding. *Gen Dent*. 2014;62(1):28–31.
12. Meijering Ac, Creugers Nhj, Roeters Fjm, Mulder J. Survival Of Three Types Of Veneer Restorations In A Clinical Trial: A 2.5-Year Interim Evaluation. *J Dent*. 1998 Sep 1;26(7):563–8.
13. Pini Nip, Sundfeld-Neto D, Aguiar Fhb, Sundfeld Rh, Martins Lrm, Lovadino Jr, Et Al. Enamel Microabrasion: An Overview Of Clinical And Scientific Considerations. *World Journal Of Clinical Cases: Wjcc*. 2015;3(1):34.
14. Janani K, Palanivelu A, Sandhya R. Diagnostic Accuracy Of Dental Pulse Oximeter With Customized Sensor Holder, Thermal Test And Electric Pulp Test For The Evaluation Of Pulp Vitality: An In Vivo Study. *Brazilian Dental Science*. 2020;23(1):8.
15. McLaren Ea, Whiteman Yy. Ceramics: Rationale For Material Selection. *Compend Contin Educ Dent*. 2010;31(9):666–8.
16. Vanini L, Mangani Fm. Determination And Communication Of Color Using The Five Color Dimensions Of Teeth. *Pract Periodontics Aesthet Dent*. 2001;13(1):19–26.
17. Dietschi D, Non-Metallici Srra. *Attuali Concetti Per Il Trattamento Estetico Dei Denti Posteriori*. Milano: Scienza Tecnica Dentistica Ed Int. 1997;
18. Vanini L, De Fs, Tammaro S. Indirect Composite Restorations In The Anterior Region: A Predictable Technique For Complex Cases. *Pract Periodontics Aesthet Dent*. 1997;9(7):795–802.
19. Caleffi A, Berardi D. Veneers In Porcellana Mordenzata: Esperienze Cliniche E Nuove Metodiche: Dalle Porcellane Tradizionali Alle Ceramiche Integrali A Pressione. Resch; 1994.
20. Hafeez N, Others. Accessory Foramen In The Middle Cranial Fossa. *Research Journal Of Pharmacy And Technology*. 2016;9(11):1880.
21. Krishnan Rp, Ramani P, Sherlin Hj, Sukumaran G, Ramasubramanian A, Jayaraj G, Et Al. Surgical Specimen Handover From Operation Theater To Laboratory: A Survey. *Ann Maxillofac Surg*. 2018 Jul;8(2):234–8.
22. Somasundaram S, Ravi K, Rajapandian K, Gurunathan D. Fluoride Content Of Bottled Drinking Water In Chennai, Tamilnadu. *J Clin Diagn Res*. 2015;9(10):Zc32.
23. Felicita As, Sumathi Felicita A. Orthodontic Extrusion Of Ellis Class Viii Fracture Of Maxillary Lateral Incisor – The Sling Shot Method [Internet]. Vol. 30, *The Saudi Dental Journal*. 2018. P. 265–9. Available From: <http://Dx.Doi.Org/10.1016/J.Sdentj.2018.05.001>
24. Kumar S, Rahman R. Knowledge, Awareness, And Practices Regarding Biomedical Waste Management Among Undergraduate Dental Students. *Asian J Pharm Clin Res*. 2017 Aug 1;10(8):341.
25. Gurunathan D, Shanmugaavel Ak. Dental Neglect Among Children In Chennai. *J Indian Soc Pedod Prev Dent*. 2016 Oct 1;34(4):364.
26. Sneha S, Others. Knowledge And Awareness Regarding Antibiotic Prophylaxis For Infective Endocarditis

- Among Undergraduate Dental Students. *Asian Journal Of Pharmaceutical And Clinical Research*. 2016;154–9.
27. Dhinesh B, Isaac Joshuaramesh Lalvani J, Parthasarathy M, Annamalai K. An Assessment On Performance, Emission And Combustion Characteristics Of Single Cylinder Diesel Engine Powered By Cymbopogon Flexuosus Biofuel. *Energy Convers Manage*. 2016 Jun 1;117:466–74.
 28. Choudhari S, Thenmozhi Ms. Occurrence And Importance Of Posterior Condylar Foramen. Laterality. 2016;8:11–43.
 29. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-Adenosine Methylation (M6a): A Promising New Molecular Target In Hypertension And Cardiovascular Diseases. *Hypertens Res*. 2020 Feb;43(2):153–4.
 30. Wu F, Zhu J, Li G, Wang J, Veeraraghavan Vp, Krishna Mohan S, Et Al. Biologically Synthesized Green Gold Nanoparticles From Siberian Ginseng Induce Growth-Inhibitory Effect On Melanoma Cells (B16). *Artif Cells Nanomed Biotechnol*. 2019 Dec;47(1):3297–305.
 31. Palati S, Ramani P, Shrelin H, Sukumaran G, Ramasubramanian A, Don Kr, Et Al. Knowledge, Attitude And Practice Survey On The Perspective Of Oral Lesions And Dental Health In Geriatric Patients Residing In Old Age Homes [Internet]. Vol. 31, *Indian Journal Of Dental Research*. 2020. P. 22. Available From: [Http://Dx.Doi.Org/10.4103/Ijdr.Ijdr_195_18](http://Dx.Doi.Org/10.4103/Ijdr.Ijdr_195_18)
 32. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis Of Silver Nanoparticles From Phenerochaete Chrysosporium (Mtcc-787) And Their Antibacterial Activity Against Human Pathogenic Bacteria. *Microb Pathog*. 2018 Apr;117:68–72.
 33. Govindaraju L, Gurunathan D. Effectiveness Of Chewable Tooth Brush In Children-A Prospective Clinical Study. *J Clin Diagn Res*. 2017;11(3):Zc31.
 34. Vijayakumar Jain S, Muthusekhar Mr, Baig Mf, Senthilnathan P, Loganathan S, Abdul Wahab Pu, Et Al. Evaluation Of Three-Dimensional Changes In Pharyngeal Airway Following Isolated Lefort One Osteotomy For The Correction Of Vertical Maxillary Excess: A Prospective Study. *J Maxillofac Oral Surg*. 2019 Mar;18(1):139–46.
 35. Coelho-De-Souza Fh, Gonçalves Ds, Sales Mp, Erhardt Mcg, Corrêa Mb, Opdam Njm, Et Al. Direct Anterior Composite Veneers In Vital And Non-Vital Teeth: A Retrospective Clinical Evaluation. *J Dent*. 2015 Nov 1;43(11):1330–6.
 36. Meijering Ac, Roeters Fjm, Mulder J, Creugers Nhj. Patients' Satisfaction With Different Types Of Veneer Restorations [Internet]. Vol. 25, *Journal Of Dentistry*. 1997. P. 493–7. Available From: [Http://Dx.Doi.Org/10.1016/S0300-5712\(96\)00067-X](http://Dx.Doi.Org/10.1016/S0300-5712(96)00067-X)
 37. Burke Fjt, Lucarotti Psk. Ten-Year Outcome Of Porcelain Laminate Veneers Placed Within The General Dental Services In England And Wales. *J Dent*. 2009 Jan 1;37(1):31–8.
 38. Burke Fjt, Lucarotti Psk, Holder Rl. Outcome Of Direct Restorations Placed Within The General Dental Services In England And Wales (Part 2): Variation By Patients' Characteristics. *J Dent*. 2005 Nov 1;33(10):817–26.
 39. Dunne Sm, Millar Bj. A Longitudinal Study Of The Clinical Performance Of Porcelain Veneers. *Br Dent J*. 1993 Nov 1;175(9):317–21.
 40. Vijayashree Priyadharsini J. In Silico Validation Of The Non-Antibiotic Drugs Acetaminophen And Ibuprofen As Antibacterial Agents Against Red Complex Pathogens. *J Periodontol*. 2019 Dec;90(12):1441–8.
 41. Pc J, Marimuthu T, Devadoss P. Prevalence And Measurement Of Anterior Loop Of The Mandibular Canal

Using Cbct: A Cross Sectional Study. Clin Implant Dent Relat Res [Internet]. 2018; Available From: <https://Europepmc.Org/Article/Med/29624863>

42. Ramesh A, Varghese S, Jayakumar Nd, Malaiappan S. Comparative Estimation Of Sulfiredoxin Levels Between Chronic Periodontitis And Healthy Patients - A Case-Control Study. J Periodontol. 2018 Oct;89(10):1241–8.
43. Ramadurai N, Gurunathan D, Samuel Av, Subramanian E, Rodrigues Sjl. Effectiveness Of 2% Articaine As An Anesthetic Agent In Children: Randomized Controlled Trial. Clin Oral Investig. 2019 Sep;23(9):3543–50.
44. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation Of Salivary Metabolomics In Oral Leukoplakia And Oral Squamous Cell Carcinoma. J Oral Pathol Med. 2019 Apr;48(4):299–306.
45. Ezhilarasan D, Apoorva Vs, Ashok Vardhan N. Syzygium Cumini Extract Induced Reactive Oxygen Species-Mediated Apoptosis In Human Oral Squamous Carcinoma Cells. J Oral Pathol Med. 2019 Feb;48(2):115–21.
46. Mathew Mg, Samuel Sr, Soni Aj, Roopa Kb. Evaluation Of Adhesion Of Streptococcus Mutans, Plaque Accumulation On Zirconia And Stainless Steel Crowns, And Surrounding Gingival Inflammation In Primary Molars: Randomized Controlled Trial. Clin Oral Investig. 2020;1–6.
47. Samuel Sr. Can 5-Year-Olds Sensibly Self-Report The Impact Of Developmental Enamel Defects On Their Quality Of Life? Int J Paediatr Dent. 2021 Mar;31(2):285–6.
48. R H, Hannah R, Ramani P, Ramanathan A, R Jm, Gheena S, Et Al. Cyp2 C9 Polymorphism Among Patients With Oral Squamous Cell Carcinoma And Its Role In Altering The Metabolism Of Benzo[A]Pyrene [Internet]. Vol. 130, Oral Surgery, Oral Medicine, Oral Pathology And Oral Radiology. 2020. P. 306–12. Available From: <http://Dx.Doi.Org/10.1016/J.Oooo.2020.06.021>
49. Chandrasekar R, Chandrasekhar S, Sundari Kks, Ravi P. Development And Validation Of A Formula For Objective Assessment Of Cervical Vertebral Bone Age. Prog Orthod. 2020 Oct 12;21(1):38.
50. Vijayashree Priyadharsini J, Smiline Girija As, Paramasivam A. In Silico Analysis Of Virulence Genes In An Emerging Dental Pathogen A. Baumannii And Related Species. Arch Oral Biol. 2018 Oct;94:93–8.

List Of Tables And Figures:

Table/Figures	Title	Page Number
Table 1	Frequency Table Showing The Age Distribution Of Patients Who Had Undergone Veneering Treatment	16
Table 2	Frequency Table Showing The Gender Distribution Of Patients Who Had Undergone Veneering Treatment	17
Table 3	Frequency Table Showing The Reason For Undergoing Veneering Treatment	18
Figure 1	Bar Chart Showing Association Between Age And Reason For Veneering Treatment	19
Figure 2	Bar Chart Showing Association Between Gender And Reason For Veneering Treatment	20

Age			Number Of Patients	Percentage
Valid	Below Years	20	18	11.5
	20 To Years	40	111	71.2
	40 To Years	60	9	5.8
	Above Years	60	18	11.5
	Total		156	100.0

Table 1 Shows The Age Distribution Of Patients Who Had Undergone Veneering Treatment. 11.5% Of The Patients Belonged To The Age Group Of Below 20 Years, 71.2% Of The Patients Belonged To The Age Group Between 20 To 40 Years, 5.8% Belonged To The Age Group Between 40 To 60 Years And 11.5% Of The Patients Belonged To The Age Group Of Above 60 Years.

Gender		Number Of Patients	Percentage
Valid	Female	59	37.8

Male	97	62.2
Total	156	100.0

Table 2 Shows The Gender Distribution Of Patients Who Had Undergone Veneering Treatment. 37.8% Of The Patients Were Females And 62.2% Of The Patients Were Males.

Reason		Number Of Patients	Percentage
Valid	Discolouration	64	41.0
	Discolouration + Spacing	16	10.3
	Spacing	76	48.7
	Total	156	100.0

Table 3 Shows The Reason For Undergoing Veneering Treatment. 41% Of The Patients Reported With The Complaint Of Discolouration, 48.7% Of The Patients Reported With The Complaint Of Spacing And 10.3% Of The Patients Reported With The Complaint Of Both Spacing And Discolouration.




Figure 1 Shows The Association Between Age And Reason For Veneering Treatment. X Axis Represents The Age Group And Y Axis Represents The Number Of Teeth. Blue Colour Denotes Discolouration, Green Colour Denotes Discolouration And Spacing And Beige Colour Denotes Spacing. In The Age Group Of Below 20 Years, 5.13% Reported With A Complaint Of Discolouration And 6.41% Of The Patients Reported With A Complaint Of Spacing. In The Age Group Between 20 To 40 Years, 24.36% Reported With A Complaint Of Discolouration, 36.54% Reported With A Complaint Of Spacing And 10.26% Of The Patients Reported With Complaints Of Both Spacing And Discolouration. In The Age Group Between 40-60 Years, Spacing Was The Only Complaint (5.77%) And In The Age Group Of Above 60 Years, Discolouration Was The Only Complaint. (11.54%) Overall, Spacing Was The Most Common Reason In The Age Group Between 20-40 Years. Chi-Square Test Was Done And The Association Was Found To Be Statistically Significant. Pearson's Chi-Square Value = 40.696, P Value 0.00 (<0.05), Statistically Significant.




Figure 2 Shows The Association Between Gender And Reason For Veneering Treatment. X Axis Represents The Gender And Y Axis Represents The Number Of Teeth. Blue Colour Denotes Discolouration, Green Colour Denotes Discoloration And Spacing And Beige Colour Denotes Spacing. Among Females, 10.26% Of The Patients Reported With A Complaint Of Discolouration And 27.56% Of The Patients Reported With A Complaint Of Spacing. Among Males, 30.77% Reported With A Complaint Of Discoloration, 21.15% Reported With A Complaint Of Spacing And 10.26% Reported With Complaints Of Both Spacing And Discoloration. Overall, Spacing Was The Most Common Reason Among Females And Discolouration Was The Most Common Reason Among Males. Chi-Square Test Was Done And The Association Was Found To Be Statistically Significant. Pearson's Chi-Square Value = 25.577, Df = 2, P Value 0.00 (<0.05) Hence Statistically Significant.